



Mr. Danny Mitry  
Consultant Ophthalmic Surgeon

## CATARACT SURGERY LEAFLET

### THE DAY OF YOUR OPERATION

- Please arrive at the Day Surgery Unit at the time indicated on your appointment letter, which you should bring with you. This is not the time of your procedure as some preparation is necessary once you have arrived.
- Please bring all your medication with you should the nurse/anaesthetist wish to review it.

### THE FIRST 24 HOURS AFTER SURGERY

- You should take great care of yourself and have a gentle day. Take particular care when stepping off pavements and going up and down stairs.
- After the surgery you will have an eye shield over the eye. These should be left on overnight. Please remove the following morning by 8 am. Clean your lids with cool boiled water and tissue paper. Instil your eye drops.
- The eye shield should be worn at night for one week after the operation to protect the eye during sleep. Stick the shield down with sellotape. The shield should be washed and dried.

### AFTERCARE

- A follow up out-patient appointment will be arranged 1-2 weeks after your operation. You may be given an appointment sooner than this if required
- You can use your old glasses after the operation, but don't expect them to give you perfect vision. You can get new glasses 4-5 weeks after the surgery.
- Your eye should not be painful but slight discomfort may be felt. Make sure you have some simple pain relief tablets at home.

### DROPS\

Please use your medication exactly as instructed. Start on the day of treatment.

**Maxidex**– steroid drop:

1 drop to the operated eye four times a day for 2 weeks, then twice a day for 2 weeks (4



weeks in total)

### **Chloramphenicol - antibiotic:**

One drop four times a day for 1 week.

Mr. Danny Mitry  
Consultant Ophthalmic Surgeon

### **How to instil your eye drops**

1. Thoroughly wash your hands
2. Stand in front of a mirror, pull the skin of your bottom eyelid down to form a pocket
3. Being careful not to touch your eye, squeeze one drop into the bottom of the eyelid

### **DO NOT**

- Do not rub or put pressure on your eyes for at least 1 week after surgery
- Do not squeeze your eyes for at least 1 day after surgery
- Do not drive until your vision has cleared and you feel comfortable to do so
- Do not wear make – up for one week
- Do not wet your eyes for one week
- Do not shower for two days
- Do not swim for at least 4 weeks then use goggles for 1 month
- Do not go back to heavy work or contact sports for at least 1 month after the operation, and when you do begin, start gradually.
- Do not get soap in your eyes when you wash your hair. Ensure this by leaning back, or by getting someone to do it for you.

### **DO**

- Do take your prescribed eye drops as instructed (see below)
- Do wear your eye shield at night for the first night
- Do wear sunglasses outside for the first week after surgery
- Do avoid smoky and dusty atmospheres
- Do avoid air travel for at least a week following your surgery
- Do keep all your follow up appointments

### **WHEN TO SEEK DOCTOR'S ADVICE**

- Any severe pain not relieved by ordinary pain tablets.
- Swelling of the eye and surrounding area.
- A sticky eye or one that becomes redder with time.
- Vision that gets worse rather than better.
- Raised temperature.
- 

**If you have any further questions please contact the office:**

**E: [secretary@mitryvision.com](mailto:secretary@mitryvision.com)**

**T: 02038268370**